

**APPLICATION FOR  
UNITED STATES PATENT**

**IN THE  
UNITED STATES PATENT AND TRADEMARK OFFICE**

(Attorney Docket No. 47508.577)

**Title:**

**MODULATION OF IMMUNOSTIMULATORY ACTIVITY OF  
IMMUNOSTIMULATORY OLIGONUCLEOTIDE ANALOGS BY POSITIONAL  
CHEMICAL CHANGES**

**Inventors:**

**Ekambar R. Kandimalla, Qiuyan Zhao, Dong Yu, Sudhir Agrawal**

**Assignee:**

**Hybridon, Inc.**

**CERTIFICATE OF MAILING (37 C.F.R. § 1.10)**

I hereby certify that this correspondence is being deposited with the United States Postal Service as "Express Mail Post Office to Addressee" Service under 37 CFR 1.110 on September 26, 2001 and is addressed to: Box Patent Application, Commissioner for Patents, Washington, D.C. 20231.

EL521665415US  
Express Mail Label No.

By: Jo-Ann Bergantino  
Jo-Ann Bergantino

**MODULATION OF IMMUNOSTIMULATORY ACTIVITY OF  
IMMUNOSTIMULATORY OLIGONUCLEOTIDE ANALOGS BY POSITIONAL  
CHEMICAL CHANGES**

(Attorney Docket No. 47508.577)

5

**CROSS REFERENCE TO RELATED APPLICATIONS**

This application is a continuation-in-part of U.S. patent application serial no. 09/712,898, filed on November 15, 2000. This application also claims priority from U.S. provisional patent application serial nos. 60/235,452 and 60/235,453, both filed on September 26, 2000. Each of the patent applications listed above is hereby incorporated by reference in its entirety.

**BACKGROUND OF THE INVENTION**

Field of the invention

The invention relates to the therapeutic use of oligonucleotides or oligonucleotide analogs as immunostimulatory agents in immunotherapy applications.

Summary of the related art

Oligonucleotides have become indispensable tools in modern molecular biology, being used in a wide variety of techniques, ranging from diagnostic probing methods to PCR to antisense inhibition of gene expression and immunotherapy applications. This widespread use of oligonucleotides has led to an increasing demand for rapid, inexpensive and efficient methods for synthesizing oligonucleotides.

The synthesis of oligonucleotides for antisense and diagnostic applications can now be routinely accomplished. See e.g., *Methods in Molecular Biology, Vol 20: Protocols for Oligonucleotides and Analogs* pp. 165-189 (S. Agrawal, Ed., Humana Press, 1993); *Oligonucleotides and Analogues: A Practical Approach*, pp. 87-108 (F. Eckstein, Ed., 1991); and Uhlmann and Peyman, *supra*. Agrawal and Iyer, *Curr. Op. in Biotech.* 6: 12 (1995); and *Antisense Research and Applications* (Crooke and Lebleu, Eds., CRC Press, Boca

Raton, 1993). Early synthetic approaches included phosphodiester and phosphotriester chemistries. Khorana et al., *J. Molec. Biol.* 72: 209 (1972) discloses phosphodiester chemistry for oligonucleotide synthesis. Reese, *Tetrahedron Lett.* 34: 3143-3179 (1978), discloses phosphotriester chemistry for synthesis of oligonucleotides and polynucleotides. These early approaches have largely given way to the more efficient phosphoramidite and H-phosphonate approaches to synthesis. Beaucage and Caruthers, *Tetrahedron Lett.* 22: 1859-1862 (1981), discloses the use of deoxynucleoside phosphoramidites in polynucleotide synthesis. Agrawal and Zamecnik, U.S. Patent No. 5,149,798 (1992), discloses optimized synthesis of oligonucleotides by the H-phosphonate approach.

Both of these modern approaches have been used to synthesize oligonucleotides having a variety of modified internucleotide linkages. Agrawal and Goodchild, *Tetrahedron Lett.* 28: 3539-3542 (1987), teaches synthesis of oligonucleotide methylphosphonates using phosphoramidite chemistry. Connolly et al., *Biochemistry* 23: 3443 (1984), discloses synthesis of oligonucleotide phosphorothioates using phosphoramidite chemistry. Jager et al., *Biochemistry* 27: 7237 (1988), discloses synthesis of oligonucleotide phosphoramidates using phosphoramidite chemistry. Agrawal et al., *Proc. Natl. Acad. Sci. USA* 85: 7079-7083 (1988), discloses synthesis of oligonucleotide phosphoramidates and phosphorothioates using H-phosphonate chemistry.

More recently, several researchers have demonstrated the validity of the use of oligonucleotides as immunostimulatory agents in immunotherapy applications. The observation that phosphodiester and phosphorothioate oligonucleotides can induce immune stimulation has created interest in developing this side effect as a therapeutic tool. These efforts have focused on phosphorothioate oligonucleotides containing the dinucleotide CpG.

Kuramoto et al., *Jpn. J. Cancer Res.* 83: 1128-1131 (1992) teaches that phosphodiester oligonucleotides containing a palindrome that includes a CpG dinucleotide can induce interferon-alpha and gamma synthesis and enhance natural killer activity. Krieg et al., *Nature* 371: 546-549 (1995) discloses that phosphorothioate

CpG-containing oligonucleotides are immunostimulatory. Liang *et al.*, *J. Clin. Invest.* 98: 1119-1129 (1996) discloses that such oligonucleotides activate human B cells.

Pisetsky, D. S.; Rich C. F., *Life Sci.* 54: 101 (1994), teaches that the immunostimulatory activity of CpG-oligos is further enhanced by the presence of phosphorothioate (PS) backbone on these oligos. Tokunaga, T.; Yamamoto, T.; Yamamoto, S. *Jap. J. Infect. Dis.* 52: 1 (1999), teaches that immunostimulatory activity of CpG-oligos is dependent on the position of CpG-motif and the sequences flanking CpG-motif. The mechanism of activation of immune stimulation by CpG-oligos has not been well understood. Yamamoto, T.; Yamamoto, S.; Kataoka, T.; Tokunaga, T., *Microbiol. Immunol.* 38: 831 (1994), however, suggests that CpG-oligos trigger immune cascade by binding to an intracellular receptor/protein, which is not characterized yet.

Several researchers have found that this ultimately triggers stress kinase pathways, activation of NF- $\kappa$ B and induction of various cytokines such as IL-6, IL-12,  $\gamma$ -IFN, and TNF- $\alpha$ . (See *e.g.*, Klinman, D. M.; Yi, A. K.; Beaucage, S. L.; Conover, J.; Krieg, A. M., *Proc. Natl. Acad. Sci. U. S. A.* 93: 2879 (1996); Sparwasser, T.; Miethke, T.; Lipford, G. B.; Erdmann, A.; Haecker, H.; Heeg, K.; Wagner, H., *Eur. J. Immunol.* 27: 1671 (1997); Lipford, G. B.; Sparwasser, T.; Bauer, M.; Zimmermann, S.; Koch, E. S.; Heeg, K.; Wagner, H. *Eur. J., Immunol.* 27: 3420 (1997); Sparwasser, T.; Koch, E. S.; Vabulas, R. M.; Lipford, G. B.; Heeg, K.; Ellart, J. W.; Wagner, H., *Eur. J. Immunol.* 28: 2045 (1998); and Zhao, Q.; Temsamani, J.; Zhou, R. Z.; Agrawal, S. *Antisense Nucleic Acid Drug Dev.* 7: 495 (1997).)

The use of CpG-PS-oligos as antitumor, antiviral, antibacterial and antiinflammatory agents and as adjuvants in immunotherapy has been reported. (See *e.g.*, Dunford, P. J.; Mulqueen, M. J.; Agrawal, S. *Antisense 97: Targeting the Molecular Basis of Disease*, (Nature Biotechnology) Conference abstract, 1997, pp 40; Agrawal, S.; Kandimalla E. R. *Mol. Med. Today* 6: 72 (2000); Chu. R. S.; Targoni, O. S.; Krieg, A. M.; Lehmann, P. V.; Harding, C. V. *J. Exp. Med.* 186: 1623 (1997); Zimmermann, S.; Egeter, O.; Hausmann, S.; Lipford, G. B.; Rocken, M.; Wagner, H.; Heeg, K. *J. Immunol.* 160: 3627 (1998).) Moldoveanu *et al.*, *Vaccine* 16: 1216-124 (1998) teaches that CpG-containing

phosphorothioate oligonucleotides enhance immune response against influenza virus. McCluskie and Davis, *J. Immunol.* 161: 4463-4466 (1998) teaches that CpG-containing oligonucleotides act as potent adjuvants, enhancing immune response against hepatitis B surface antigen.

5 Zhao, Q.; Temsamani, J.; Idarola, P.; Jiang, Z.; Agrawal, S. *Biochem. Pharmacol.* 51: 173 (1996), teaches that replacement of deoxynucleosides in a CpG-motif with 2'-O-methylribonucleosides suppresses immunostimulatory activity, suggesting that a rigid C3'-endo conformation induced by 2'-O-methyl modification does not allow proper recognition and/or interaction of CpG-motif with the proteins involved in the  
10 immunostimulatory pathway. This reference further teaches that substitution of a methyl group for an unbridged oxygen on the phosphate group between C and G of a CpG-motif suppresses immune stimulatory activity, suggesting that negative charge on phosphate group is essential for protein recognition and interaction.

15 Zhao, Q.; Yu, D.; Agrawal, S. *Bioorg. Med. Chem. Lett.* 9: 3453 (1999), teaches that substitution of one or two 2'-deoxynucleosides adjacent to CpG-motif with 2'- or 3'-O-methylribonucleosides on the 5'-side causes a decrease in immunostimulatory activity, while the same substitutions have insignificant effect when they were placed on the 3'-side of the CpG-motif. However, Zhao, Q.; Yu, D.; Agrawal, S. *Bioorg. Med. Chem. Lett.* 10: 1051 (2000), teaches that the substitution of a deoxynucleoside two or three  
20 nucleosides away from the CpG-motif on the 5'-side with one or two 2'-O-methoxyethyl- or 2'- or 3'-O-methylribonucleosides results in a significant increase in immunostimulatory activity.

The precise structural requirements and specific functional groups of CpG-motif necessary for the recognition of protein/receptor factor that is responsible for immune  
25 stimulation have not yet been studied in detail. There is, therefore, a need for new immunostimulatory motifs which may provide improved immunostimulatory activity.

## BRIEF SUMMARY OF THE INVENTION

The invention provides methods for enhancing the immune response caused by immunostimulatory oligonucleotide compounds. The methods according to the invention enable increasing the immunostimulatory effect for immunotherapy applications. Thus, the invention further provides methods for making and using such oligonucleotide compounds.

The present inventors have surprisingly discovered that positional modification of immunostimulatory oligonucleotides dramatically affects their immunostimulatory capabilities. In particular, modifications in the immunostimulatory domain and/or the potentiation domain enhance the immunostimulatory effect in a reproducible and predictable manner.

In a first aspect, the invention provides immunostimulatory oligonucleotide compounds comprising an immunostimulatory domain and, optionally, one or more potentiation domains. In some embodiments, the immunostimulatory domain comprises a dinucleotide analog that includes a non-naturally occurring pyrimidine base. In some embodiments, the immunostimulatory domain and/or the potentiation domain include an immunostimulatory moiety at a specified position, as described hereinbelow. In some embodiments, the immunostimulatory oligonucleotide comprises a 3'-3' linkage. In one embodiment, such 3'-3' linked oligonucleotides have two accessible 5'-ends.

In a second aspect, the invention provides methods for modulating the immunostimulatory effect of an immunostimulatory oligonucleotide compound. In some embodiments, the method comprises introducing into the immunostimulatory domain a dinucleotide analog that includes a non-naturally occurring pyrimidine base. In some embodiments, the method comprises introducing into the immunostimulatory domain and/or potentiation domain an immunostimulatory moiety at a specified position, as described hereinbelow. In some embodiments, the method comprises introducing into the oligonucleotide a 3'-3' linkage.

In a third aspect, the invention provides methods for generating an immune response in a patient, such methods comprising administering to the patient an immunostimulatory oligonucleotide compound according to the invention.

5 In a fourth aspect, the invention provides methods for therapeutically treating a patient having disease caused by a pathogen, such methods comprising administering to the patient an immunostimulatory oligonucleotide compound according to the invention.

10 In a fifth aspect, the invention provides methods for treating a cancer patient, such methods comprising administering to the patient an immunostimulatory oligonucleotide compound according to the invention.

15 In a sixth aspect, the invention provides methods for treating autoimmune disorders, such as autoimmune asthma, such methods comprising administering to the patient an oligonucleotide analog immunostimulatory compound according to the invention. Administration is carried out as described for the third aspect of the invention.

20 In a seventh aspect, the invention provides methods for treating airway inflammation or allergies, such methods comprising administering to the patient an oligonucleotide analog immunostimulatory compound according to the invention. Administration is carried out as described for the third aspect of the invention.

## BRIEF DESCRIPTION OF THE FIGURES

Sub  
A'  
Figure 1 shows results of proliferation assays using oligonucleotides having 1',2'-dideoxyribose substitutions at various positions.

5 Figure 2 shows results of spleen weight assays using oligonucleotides having 1',2'-dideoxyribose substitutions at various positions.

Figure 3 shows results of proliferation assays using different oligonucleotides having 1',2'-dideoxyribose substitutions at various positions.

Figure 4 shows results of spleen weight assays using different oligonucleotides having 1',2'-dideoxyribose substitutions at various positions.

10 Figure 5 shows results of proliferation assays using oligonucleotides having C3-linker substitutions at various positions.

Figure 6 shows results of spleen weight assays using oligonucleotides having C3-linker substitutions at various positions.

15 Figure 7 shows results of proliferation assays using oligonucleotides having Spacer 9 or Spacer 18 substitutions at various positions.

Figure 8 shows results of spleen weight assays using oligonucleotides having Spacer 9 or Spacer 18 substitutions at various positions.

Figure 9 shows results of proliferation assays using oligonucleotides having amino-linker substitutions at various positions.

20 Figure 10 shows results of spleen weight assays using oligonucleotides having amino-linker substitutions at various positions.

Figure 11 shows results of proliferation assays using oligonucleotides having 3'-deoxynucleoside substitutions at various positions.

25 Figure 12 shows results of spleen weight assays using oligonucleotides having 3'-deoxynucleoside substitutions at various positions.



Sub  
A1  
Figure 13 shows results of proliferation assays using oligonucleotides having methylphosphonate substitutions at various positions.

Figure 14 shows results of spleen weight assays using oligonucleotides having methylphosphonate substitutions at various positions.

5 Figure 15 shows results of proliferation assays using oligonucleotides having 2'-O-methylribonucleoside or 2'-O-methoxyethyl substitutions at various positions.

Figure 16 shows results of spleen weight assays using oligonucleotides having 2'-O-methylribonucleoside or 2'-O-methoxyethyl substitutions at various positions.

Figure 17 shows results of proliferation assays using oligonucleotides having 5'-3', 5'-5', or 3'-3' linkage substitutions at various positions.

Figure 18 shows results of spleen weight assays using oligonucleotides having  $\beta$ -L-deoxynucleotide substitutions at various positions.

Figure 19 shows results of spleen weight assays using oligonucleotides having 2'-O-propargyl substitutions at various positions.

Figure 20 shows results of spleen weight assays using oligonucleotides having various substitutions at various positions.

Figure 21 shows results of spleen weight assays using oligonucleotides having 7-deazaguanine substitution within the immunostimulatory dinucleotide.

20 Figure 22 shows results of proliferation assays using oligonucleotides having 6-thioguanine substitution within the immunostimulatory dinucleotide.

Figure 23 shows results of spleen weight assays using oligonucleotides having 5-hydroxycytosine or N4-ethylcytosine substitution within the immunostimulatory dinucleotide.

25 Figure 24 shows results of spleen weight assays using oligonucleotides having 5-hydroxycytosine or N4-ethylcytosine substitution within the immunostimulatory dinucleotide.

Figure 25 shows results of proliferation assays using oligonucleotides having arabinofuranosylcytosine (aracytidine; Ara-C) substitution within the immunostimulatory dinucleotide.

Figure 26 shows results of spleen weight assays using oligonucleotides having 4-thiouracil substitution within the immunostimulatory dinucleotide.

Figure 27 shows the chemical structure of a CpG-motif, showing functional groups on cytosine that serve as hydrogen bond acceptor and hydrogen bond donor groups.

Figure 28 shows the chemical structures of cytosine (1) and cytosine analogs (2-7). In the nucleosides cytidine, deoxycytidine, and related analogs, the substituent R is ribose or 2'-deoxyribose.

10  
09/26/01  
14:29:50  
T0220

## DETAILED DESCRIPTION

The invention relates to the therapeutic use of oligonucleotides and oligonucleotide analogs as immunostimulatory agents for immunotherapy applications. The patents and publications cited herein reflect the level of knowledge in the field and are hereby incorporated by reference in their entirety. In the event of conflict between any teaching of any reference cited herein and the present specification, the latter shall prevail, for purposes of the invention.

The invention provides methods for enhancing the immune response caused by immunostimulatory oligonucleotide compounds for immunotherapy applications. Thus, the invention further provides compounds having optimal levels of immunostimulatory effect for immunotherapy and methods for making and using such oligonucleotide compounds.

The present inventors have surprisingly discovered that positional chemical modifications introduced in immunostimulatory oligonucleotides dramatically affect their immunostimulatory capabilities. In particular, modifications in the immunostimulatory domain and/or the potentiation domain can enhance the immunostimulatory effect in a reproducible manner for desired applications.

In a first aspect, the invention provides immunostimulatory oligonucleotide compounds comprising an immunostimulatory domain and, optionally, one or more potentiation domains. In certain preferred embodiments, the immunostimulatory domain comprises a dinucleotide analog that includes a non-natural pyrimidine nucleoside.

For purposes of all aspects of the invention, the term "oligonucleotide" includes polymers of two or more deoxyribonucleosides, or any modified nucleoside, including 2'- or 3'-substituted nucleosides, 2'- or 3'-O-substituted ribonucleosides, deazanucleosides, or any combination thereof. Such monomers may be coupled to each other by any of the numerous known internucleoside linkages. In certain preferred embodiments, these internucleoside linkages may be phosphodiester, phosphotriester,

phosphorothioate, phosphorodithioate, or phosphoramidate linkages, including 3'-5', 2'-5', 3'-3', and 5'-5' linkages of any of the foregoing, or combinations thereof. The term oligonucleotide also encompasses such polymers having chemically modified bases or sugars and/or having additional substituents, including without limitation lipophilic groups, intercalating agents, diamines and adamantane. The term oligonucleotide also encompasses peptide nucleic acids (PNA), peptide nucleic acids with phosphate groups (PHONA), locked nucleic acids (LNA), morpholinonucleic acids, and oligonucleotides comprising non-pentose sugar (e.g. hexose) or abasic sugar backbones or backbone sections, as well as oligonucleotides that include backbone sections with non-sugar linker or spacer groups, as further described hereinbelow.


For purposes of the invention the terms "2'-substituted" and "3'-substituted" mean (respectively) substitution of the 2' (or 3') position of the pentose moiety with a halogen (preferably Cl, Br, or F), or an -O-lower alkyl group containing 1-6 saturated or unsaturated carbon atoms, or with an -O-aryl or allyl group having 2-6 carbon atoms, wherein such alkyl, aryl or allyl group may be unsubstituted or may be substituted, e.g., with halo, hydroxy, trifluoromethyl, cyano, nitro, acyl, acyloxy, alkoxy, carboxyl, carbalkoxy, or amino groups; or such 2' substitution may be with a hydroxy group (to produce a ribonucleoside) or an amino group, but not with a 2' (or 3') H group.

For purposes of the invention, the term "immunostimulatory oligonucleotide compound" means a compound comprising an immunostimulatory dinucleotide, without which the compound would not have an immunostimulatory effect. An "immunostimulatory dinucleotide" is a dinucleotide having the formula 5'-pyrimidine-purine-3', wherein "pyrimidine" is a natural or non-natural pyrimidine nucleoside and "purine" is a natural or non-natural purine nucleoside. One such immunostimulatory dinucleotide is CpG. The terms "CpG" and "CpG dinucleotide" refer to the dinucleotide 5'-deoxycytidine-deoxyguanosine-3', wherein p is an internucleotide linkage, preferably selected from the group consisting of phosphodiester, phosphorothioate, and phosphorodithioate.

For purposes of the invention, a "dinucleotide analog" is an immunostimulatory dinucleotide as described above, wherein either or both of the pyrimidine and purine nucleosides is a non-natural nucleoside. A "non-natural" nucleoside is one that includes a non-naturally occurring base and/or a non-naturally occurring sugar moiety. For  
5 purposes of the invention, a base is considered to be non-natural if it is not selected from the group consisting of thymine, guanine, cytosine, adenine, and uracil. The terms "C\*pG" and "CpG\*" refer to immunostimulatory dinucleotide analogs comprising a cytidine analog (non-natural pyrimidine nucleoside) or a guanosine analog (non-natural purine nucleoside), respectively.

10 Figure 27 shows the chemical structure of a CpG-motif, showing the functional groups on cytosine that serve as hydrogen bond acceptor and hydrogen bond donor groups. Cytosine has two hydrogen bond acceptor groups at positions 2 (keto-oxygen) and 3 (nitrogen), and a hydrogen bond donor group at the 4-position (amino group). These groups can serve as potential recognizing and interacting groups with receptors that are responsible for immune stimulation. Figure 28 shows cytosine analogs that are  
15 isostructural with natural cytosine, including 5-methyl-deoxycytosine (2), 5-methyl-deoxyisocytosine (3), 5-hydroxy-deoxycytosine (4), deoxyuridine (5), N4-ethyl-deoxycytosine (6), and deoxy-P-base (7).

In one embodiment, therefore, the immunostimulatory dinucleotide comprises a  
20 pyrimidine nucleoside of structure (I):




(I)

wherein D is a hydrogen bond donor, D' is selected from the group consisting of hydrogen, hydrogen bond donor, hydrogen bond acceptor, hydrophilic group,  
25 hydrophobic group, electron withdrawing group and electron donating group, A is a

hydrogen bond acceptor, X is carbon or nitrogen, and S is a pentose or hexose sugar ring linked to the pyrimidine base. In some embodiments, the pyrimidine nucleoside is a non-natural pyrimidine nucleoside, i.e., the compound of structure (I) is not cytidine or deoxycytidine.

5 In some embodiments, the base moiety in (I) is a non-naturally occurring pyrimidine base. Examples of preferred non-naturally occurring pyrimidine bases include, without limitation, 5-hydroxycytosine, 5-hydroxymethylcytosine, N4-alkylcytosine, preferably N4-ethylcytosine, and 4-thiouracil. In some embodiments, the sugar moiety S in (I) is a non-naturally occurring sugar moiety. For purposes of the present invention, a "naturally occurring sugar moiety" is ribose or 2'-deoxyribose, and a "non-naturally occurring sugar moiety" is any sugar other than ribose or 2'-deoxyribose that can be used in the backbone for an oligonucleotide.. Arabinose and arabinose derivatives are examples of a preferred non-naturally occurring sugar moieties.

15 Immunostimulatory domains according to the invention may include immunostimulatory moieties on one or both sides of the immunostimulatory natural dinucleotide or non-natural dinucleotide analog. For example, an immunostimulatory domain could be depicted as



20 wherein Y represents cytidine or a non-natural pyrimidine nucleoside analog, Z represents guanosine or a non-natural purine nucleoside analog, and each X independently represents a nucleoside or an immunostimulatory moiety according to the invention. An "immunostimulatory moiety" is a chemical structure at a particular position within the immunostimulatory domain or the potentiation domain that causes  
25 the immunostimulatory oligonucleotide to be more immunostimulatory than it would be in the absence of the immunostimulatory moiety.

Preferred immunostimulatory moieties include modifications in the phosphate backbones including without limitation methylphosphonates, methylphosphonothioates phosphotriesters, phosphothiotriesters phosphorothioates, phosphorodithioates, triester prodrugs, sulfones, sulfonamides, sulfamates, formacetal, N-methylhydroxylamine, carbonate, carbamate, boranophosphonate, phosphoramidates, especially primary amino-phosphoramidates, N3 phosphoramidates and N5 phosphoramidates, and stereospecific linkages (e.g., (R)- or (S)-phosphorothioate, alkylphosphonate, or phosphotriester linkages). Preferred immunostimulatory moieties according to the invention further include nucleosides having sugar modifications, including without limitation 2'-substituted pentose sugars including without limitation 2'-O-methylribose, 2'-O-methoxyethylribose, 2'-O-propargylribose, and 2'-deoxy-2'-fluororibose; 3'-substituted pentose sugars, including without limitation 3'-O-methylribose; 1',2'-dideoxyribose; hexose sugars, including without limitation arabinose, 1'-methyларabınose, 3'-hydroxymethyларabınose, 4'-hydroxymethyларabınose, and 2'-substituted arabinose sugars; and alpha-anomers.


Preferred immunostimulatory moieties according to the invention further include oligonucleotides having other carbohydrate backbone modifications and replacements, including peptide nucleic acids (PNA), peptide nucleic acids with phosphate groups (PHONA), locked nucleic acids (LNA), morpholinonucleic acids, and oligonucleotides having backbone sections with alkyl linkers or amino linkers. The alkyl linker may be branched or unbranched, substituted or unsubstituted, and chirally pure or a racemic mixture. Most preferably, such alkyl linkers have from about 2 to about 18 carbon atoms. In some preferred embodiments such alkyl linkers have from about 3 to about 9 carbon atoms. Such alkyl linkers include polyethyleneglycol linkers  $[-O-CH_2-CH_2-]_n$  ( $n = 2-9$ ). In some preferred embodiments, such alkyl linkers may include peptides or amino acids.

Preferred immunostimulatory moieties according to the invention further include DNA isoforms, including without limitation  $\beta$ -L-deoxynucleosides and alpha-deoxynucleosides. Preferred immunostimulatory moieties according to the invention

further include nucleosides having unnatural internucleoside linkage positions, including without limitation 2'-5', 2'-2', 3'-3' and 5'-5' linkages.

Preferred immunostimulatory moieties according to the invention further include nucleosides having modified heterocyclic bases, including without limitation 5-hydroxydeoxycytidine, 5-hydroxymethyldeoxycytidine, N4-alkyldeoxycytidine, preferably N4-ethyldeoxycytidine, 4-thiouridine, 6-thiodeoxyguanosine, 7-deazaguanosine, and deoxyribonucleosides of nitropyrrole, C5-propynylpyrimidine, and diaminopurine, including without limitation 2,6-diaminopurine.

By way of specific illustration and not by way of limitation, for example, in the immunostimulatory domain described earlier



a nucleoside methylphosphonate at position X3 or X4 is an immunostimulatory moiety, a substituted or unsubstituted alkyl linker at position X1 is an immunostimulatory moiety, and a  $\beta$ -L-deoxynucleoside at position X1 is an immunostimulatory moiety. See Table 1 below for representative positions and structures of immunostimulatory moieties within the immunostimulatory domain.

Table 1

Position	TYPICAL IMMUNOSTIMULATORY MOIETIES
X1	C3-alkyl linker, 2-aminobutyl-1,3-propanediol linker (amino linker), $\beta$ -L-deoxynucleoside
X2	2-aminobutyl-1,3-propanediol linker
X3	nucleoside methylphosphonate
X4	nucleoside methylphosphonate, 2'-O-methyl-ribonucleoside

In some embodiments, the immunostimulatory oligonucleotide further comprises a potentiation domain



A "potentiation domain" is a region of an immunostimulatory oligonucleotide analog, other than the immunostimulatory domain, that causes the oligonucleotide to be more immunostimulatory if it contains the potentiation domain than the oligonucleotide would be in the absence of the potentiation domain. The potentiation domain can be upstream or downstream relative to the immunostimulatory domain. The term "upstream" is used to refer to positions on the 5' side of the immunostimulatory dinucleotide or dinucleotide analog (Y-Z). The term "downstream" is used to refer to positions on the 3' side of Y-Z.

For example, an immunostimulatory oligonucleotide analog could have the structure

5'-U9-U8-U7-U6-U5-U4-U3-U2-U1-X1-X2-Y-Z-X3-X4-N-N-N-3'

wherein U9-U1 represents an upstream potentiation domain, wherein each U independently represents the same or a different nucleoside immunostimulatory moiety, N represents any nucleoside and X1-X4, Y and Z are as before.

Alternatively, an immunostimulatory oligonucleotide analog could have the structure

5'-N-N-X1-X2-Y-Z-X3-X4-D1-D2-D3-D4-D5-D6-D7-D8-3'

wherein D1-D8 represents a downstream potentiation domain, wherein each D independently represents the same or a different nucleoside or immunostimulatory moiety, and all other symbols are as described above.

In these configurations, an immunostimulatory moiety at U6 would be six positions upstream from the immunostimulatory dinucleotide or dinucleotide analog and an immunostimulatory moiety at D4 would be four positions downstream from the immunostimulatory dinucleotide or dinucleotide analog. The term "position" is used rather than "nucleoside", because any of the U or D positions can represent an immunostimulatory moiety which may or may not be a nucleoside or nucleoside

analog. Of course, oligonucleotide analogs can be constructed having both upstream and downstream potentiation domains.

Table 2 shows representative positions and structures of immunostimulatory moieties within an immunostimulatory oligonucleotide having an upstream potentiation domain. See Figure 7 for definitions of Spacer 9 and Spacer 18 as referred to in Tables 2 and 3.

Table 2

Position	TYPICAL IMMUNOSTIMULATORY MOIETY
X2	2-aminobutyl-1,3-propanediol linker
X1	C3-linker, 2-aminobutyl-1,3-propanediol linker, $\beta$ -L-deoxy-nucleoside
U1	1',2'-dideoxyribose, C3-linker, 2'-O-methyl-ribonucleoside
U2	1',2'-dideoxyribose, C3-linker, Spacer 18, 3'-deoxynucleoside, nucleoside methylphosphonate, $\beta$ -L-deoxynucleoside, 2'-O-propargyl-ribonucleoside
U3	1',2'-dideoxyribose, C3-linker, Spacer 9, Spacer 18, nucleoside methylphosphonate, 2'-5' linkage
U2 + U3	1',2'-dideoxyribose, C3-linker, , $\beta$ -L-deoxynucleoside
U3 + U4	nucleoside methylphosphonate, 2'-O-methoxyethyl-ribonucleoside
U5 + U6	1',2'-dideoxyribose, C3-linker
X1 + U3	1',2'-dideoxyribose

Table 3 shows representative positions and structures of immunostimulatory moieties within an immunostimulatory oligonucleotide having a downstream potentiation domain.

Table 3

Position	TYPICAL IMMUNOSTIMULATORY MOIETY
X3	nucleoside methylphosphonate
X4	nucleoside methylphosphonate, 2'-O-methyl-ribonucleoside
D1	1',2'-dideoxyribose, nucleoside methylphosphonate
D2	1',2'-dideoxyribose, C3-linker, Spacer 9, Spacer 18, 2-aminobutyl-1,3-propanediol -linker, nucleoside methylphosphonate, $\beta$ -L-deoxynucleoside
D3	3'-deoxynucleoside, 2'-O-propargyl-ribonucleoside, 2'-5'-linkage
D2 + D3	1',2'-dideoxyribose, $\beta$ -L-deoxynucleoside

In another embodiment of the invention, the oligonucleotide according to the invention has one or two accessible 5' ends. The present inventors have discovered that immunostimulatory moieties in the region 5' to the immunostimulatory dinucleotide have a greater impact on immunostimulatory activity than do similar substitutions in the region 3' to the immunostimulatory dinucleotide. This observation suggests that the 5'-flanking region of CpG-PS-oligos plays an important role in immunostimulatory activity. Moreover, the inventors have discovered that compounds having two oligonucleotide units attached by way of a 3'-5' or 3'-3' linkage have greater immunostimulatory activity than do compounds in which the two oligonucleotide units are attached by way of a 5'-5' linkage. In some preferred embodiments, therefore, the immunostimulatory oligonucleotide according to the invention comprises a 3'-3' linkage. In some such embodiments, the oligonucleotides have one or two accessible 5' ends.

In a second aspect, the invention provides methods for modulating the immunostimulatory effect of an immunostimulatory oligonucleotide. In some embodiments, the method comprises introducing into the immunostimulatory domain a dinucleotide analog that includes a non-naturally occurring pyrimidine base, as described above for the first aspect of the invention. In some embodiments, the method

comprises introducing into the immunostimulatory domain and/or potentiation domain an immunostimulatory moiety at a specified position, as described above. In some embodiments, the method comprises introducing into the oligonucleotide a 3'-3' linkage.

5 For purposes of the invention, "introducing an immunostimulatory moiety" at a specified position simply means synthesizing an oligonucleotide that has an immunostimulatory moiety at the specified position. For example, "introducing an immunostimulatory moiety into position U6" simply means synthesizing an oligonucleotide that has an immunostimulatory moiety at such a position, with  
10 reference to, *e.g.*, the following structure:

5'-U9-U8-U7-U6-U5-U4-U3-U2-U1-X1-X2-Y-Z-X3-X4-D1-D2-D3-3'.

15 Preferably, the methods according to this aspect of the invention include introducing an immunostimulatory moiety at a position in the immunostimulatory domain or in an upstream or downstream potentiation domain according to the preferred substitution patterns described in Tables 1-3.

20 The methods according to this aspect of the invention can be conveniently carried out using any of the well-known synthesis techniques by simply using an appropriate immunomodulatory moiety monomer synthon in the synthesis process in an appropriate cycle to obtain the desired position. Preferred monomers include  
25 phosphoramidites, phosphotriesters and H-phosphonates. PS-oligos are readily synthesized, *e.g.*, using  $\beta$ -cyanoethylphosphoramidite chemistry on CPG solid support using appropriate phosphoramidites, deprotected as required, purified by  $C_{18}$  reverse phase HPLC, dialyzed against distilled water and lyophilized. The purity of each PS-oligo is readily determined by CGE and the molecular weight can be confirmed by  
MALDI-TOF mass spectral analysis.

In a third aspect, the invention provides methods for generating an immune response in a patient, such methods comprising administering to the patient an oligonucleotide analog immunostimulatory compound according to the invention.

In the methods according to this aspect of the invention, preferably,  
5 administration of compounds is parenteral, oral, sublingual, transdermal, topical, intranasal, intratracheal, intravaginal, or intrarectal. Administration of the therapeutic compositions can be carried out using known procedures at dosages and for periods of time effective to reduce symptoms or surrogate markers of the disease. When administered systemically, the therapeutic composition is preferably administered at a  
10 sufficient dosage to attain a blood level of oligonucleotide from about 0.001 micromolar to about 10 micromolar. For localized administration, much lower concentrations than this may be effective, and much higher concentrations may be tolerated. Preferably, a total dosage of oligonucleotide will range from about 0.1 mg oligonucleotide per patient per day to about 40 mg oligonucleotide per kg body weight per day. It may be  
15 desirable to administer simultaneously, or sequentially a therapeutically effective amount of one or more of the therapeutic compositions of the invention to an individual as a single treatment episode. In some instances, dosages below the above-defined ranges may still provide efficacy. In a preferred embodiment, after the composition of matter is administered, one or more measurement is taken of biological effects selected  
20 from the group consisting of complement activation, mitogenesis and inhibition of thrombin clot formation.

In certain preferred embodiments, compounds according to the invention are administered in combination with antibiotics, antigens, allergens, vaccines, antibodies, cytotoxic agents, antisense oligonucleotides, gene therapy vectors, DNA vaccines  
25 and/or adjuvants to enhance the specificity or magnitude of the immune response. Either the compound or the vaccine, or both may optionally be linked to an immunogenic protein, such as keyhole limpet hemocyanin, cholera toxin B subunit, or any other immunogenic carrier protein. Any of a plethora of adjuvants may be used, including, without limitation, Freund's complete adjuvant, monophosphoryl lipid A

(MPL), saponins, including QS-21, alum, and combinations thereof. Certain preferred embodiments of the methods according to the invention induce cytokines by administration of immunostimulatory oligonucleotide compounds. In certain embodiments the immunostimulatory oligonucleotide compounds are conjugated to an antigen, hapten, or vaccine. As discussed above, the present inventors have discovered that an accessible 5' end is important to the activity of certain immunostimulatory oligonucleotide compounds. Accordingly, for optimum immunostimulatory activity, the oligonucleotide preferably is conjugated to an antigen or vaccine by means of the 3'-end of oligonucleotide compound.

For purposes of this aspect "in combination with" means in the course of treating the same disease in the same patient, and includes administering the oligonucleotide and/or the vaccine and/or the adjuvant in any order, including simultaneous administration, as well as temporally spaced order of up to several days apart. Such combination treatment may also include more than a single administration of the oligonucleotide, and/or independently the vaccine, and/or independently the adjuvant. The administration of the oligonucleotide and/or vaccine and/or adjuvant may be by the same or different routes.

The method according to this aspect of the invention is useful for model studies of the immune system, and is further useful for the therapeutic treatment of human or animal disease.

In a fourth aspect, the invention provides methods for therapeutically treating a patient having disease caused by a pathogen, such methods comprising administering to the patient an oligonucleotide analog immunostimulatory compound according to the invention. Administration is carried out as described for the third aspect of the invention.

In a fifth aspect, the invention provides methods for treating a cancer patient, such methods comprising administering to the patient an oligonucleotide analog

In a sixth aspect, the invention provides methods for treating autoimmune disorders, such as autoimmune asthma, such methods comprising administering to the patient an oligonucleotide analog immunostimulatory compound according to the invention. Administration is carried out as described for the third aspect of the invention.

10  
 11  
 12  
 13  
 14  
 15  
 16  
 17  
 18  
 19  
 20  
 21  
 22  
 23  
 24  
 25  
 26  
 27  
 28  
 29  
 30  
 31  
 32  
 33  
 34  
 35  
 36  
 37  
 38  
 39  
 40  
 41  
 42  
 43  
 44  
 45  
 46  
 47  
 48  
 49  
 50  
 51  
 52  
 53  
 54  
 55  
 56  
 57  
 58  
 59  
 60  
 61  
 62  
 63  
 64  
 65  
 66  
 67  
 68  
 69  
 70  
 71  
 72  
 73  
 74  
 75  
 76  
 77  
 78  
 79  
 80  
 81  
 82  
 83  
 84  
 85  
 86  
 87  
 88  
 89  
 90  
 91  
 92  
 93  
 94  
 95  
 96  
 97  
 98  
 99  
 100  
 101  
 102  
 103  
 104  
 105  
 106  
 107  
 108  
 109  
 110  
 111  
 112  
 113  
 114  
 115  
 116  
 117  
 118  
 119  
 120  
 121  
 122  
 123  
 124  
 125  
 126  
 127  
 128  
 129  
 130  
 131  
 132  
 133  
 134  
 135  
 136  
 137  
 138  
 139  
 140  
 141  
 142  
 143  
 144  
 145  
 146  
 147  
 148  
 149  
 150  
 151  
 152  
 153  
 154  
 155  
 156  
 157  
 158  
 159  
 160  
 161  
 162  
 163  
 164  
 165  
 166  
 167  
 168  
 169  
 170  
 171  
 172  
 173  
 174  
 175  
 176  
 177  
 178  
 179  
 180  
 181  
 182  
 183  
 184  
 185  
 186  
 187  
 188  
 189  
 190  
 191  
 192  
 193  
 194  
 195  
 196  
 197  
 198  
 199  
 200  
 201  
 202  
 203  
 204  
 205  
 206  
 207  
 208  
 209  
 210  
 211  
 212  
 213  
 214  
 215  
 216  
 217  
 218  
 219  
 220  
 221  
 222  
 223  
 224  
 225  
 226  
 227  
 228  
 229  
 230  
 231  
 232  
 233  
 234  
 235  
 236  
 237  
 238  
 239  
 240  
 241  
 242  
 243  
 244  
 245  
 246  
 247  
 248  
 249  
 250  
 251  
 252  
 253  
 254  
 255  
 256  
 257  
 258  
 259  
 260  
 261  
 262  
 263  
 264  
 265  
 266  
 267  
 268  
 269  
 270  
 271  
 272  
 273  
 274  
 275  
 276  
 277  
 278  
 279  
 280  
 281  
 282  
 283  
 284  
 285  
 286  
 287  
 288  
 289  
 290  
 291  
 292  
 293  
 294  
 295  
 296  
 297  
 298  
 299  
 300  
 301  
 302  
 303  
 304  
 305  
 306  
 307  
 308  
 309  
 310  
 311  
 312  
 313  
 314  
 315  
 316  
 317  
 318  
 319  
 320  
 321  
 322  
 323  
 324  
 325  
 326  
 327  
 328  
 329  
 330  
 331  
 332  
 333  
 334  
 335  
 336  
 337  
 338  
 339  
 340  
 341  
 342  
 343  
 344  
 345  
 346  
 347  
 348  
 349  
 350  
 351  
 352  
 353  
 354  
 355  
 356  
 357  
 358  
 359  
 360  
 361  
 362  
 363  
 364  
 365  
 366  
 367  
 368  
 369  
 370  
 371  
 372  
 373  
 374  
 375  
 376  
 377  
 378  
 379  
 380  
 381  
 382  
 383  
 384  
 385  
 386  
 387  
 388  
 389  
 390  
 391  
 392  
 393  
 394  
 395  
 396  
 397  
 398  
 399  
 400  
 401  
 402  
 403  
 404  
 405  
 406  
 407  
 408  
 409  
 410  
 411  
 412  
 413  
 414  
 415  
 416  
 417  
 418  
 419  
 420  
 421  
 422  
 423  
 424  
 425  
 426  
 427  
 428  
 429  
 430  
 431  
 432  
 433  
 434  
 435  
 436  
 437  
 438  
 439  
 440  
 441  
 442  
 443  
 444  
 445  
 446  
 447  
 448  
 449  
 450  
 451  
 452  
 453  
 454  
 455  
 456  
 457  
 458  
 459  
 460  
 461  
 462  
 463  
 464  
 465  
 466  
 467  
 468  
 469  
 470  
 471  
 472  
 473  
 474  
 475  
 476  
 477  
 478  
 479  
 480  
 481  
 482  
 483  
 484  
 485  
 486  
 487  
 488  
 489  
 490  
 491  
 492  
 493  
 494  
 495  
 496  
 497  
 498  
 499  
 500  
 501  
 502  
 503  
 504  
 505  
 506  
 507  
 508  
 509  
 510  
 511  
 512  
 513  
 514  
 515  
 516  
 517  
 518  
 519  
 520  
 521  
 522  
 523  
 524  
 525  
 526  
 527  
 528  
 529  
 530  
 531  
 532

The following examples are intended to further illustrate certain preferred embodiments of the invention, and are not intended to limit the scope of the invention.

## EXAMPLES

### **Example 1: Synthesis of oligonucleotides containing immunomodulatory moieties**

Oligonucleotides were synthesized on a 1 micromolar scale using an automated DNA synthesizer (Expedite 8909, PerSeptive Biosystems, Foster City, CA). Standard deoxynucleoside phosphoramidites are obtained from PerSeptive Biosystems. 1',2'-dideoxyribose phosphoramidite, propyl-1-phosphoramidite, 2'-deoxy-5-nitroindole-ribofuranosyl phosphoramidite, 2'-deoxy-uridine phosphoramidite, 2'-deoxy-P phosphoramidite, 2'-deoxy-2-aminopurine phosphoramidite, 2'-deoxy-nebularine phosphoramidite, 2'-deoxy-7-deazaguanosine phosphoramidite, 2'-deoxy-4-thiouridine phosphoramidite, 2'-deoxy-isoguanosine phosphoramidite, 2'-deoxy-5-methylisocytosine phosphoramidite, 2'-deoxy-4-thiothymidine phosphoramidite, 2'-deoxy-K-phosphoramidite, 2'-deoxy-2-aminoadenosine phosphoramidite, 2'-deoxy-N4-ethyl-cytosine phosphoramidite, 2'-deoxy-6-thioguanosine phosphoramidite, 2'-deoxy-7-deaza-xanthosine phosphoramidite, 2'-deoxy-8-bromoguanosine phosphoramidite, 2'-deoxy-8-oxoguanosine phosphoramidite, 2'-deoxy-5-hydroxycytosine phosphoramidite, arabino-cytosine phosphoramidite and 2'-deoxy-5-propynecytosine phosphoramidite were obtained from Glen Research (Sterling, VA). 2'-Deoxy-inosine phosphoramidite were obtained from ChemGenes (Ashland, MA).

Normal coupling cycles or a coupling cycle recommended by the phosphoramidite manufacturer were used for all phosphoramidites. Beaucage reagent was used as an oxidant to obtain phosphorothioate modification. After synthesis, oligonucleotides were deprotected by incubating CPG-bound oligonucleotide with concentrated ammonium hydroxide solution for 1.5-2 hours at room temperature and then incubating the ammonium hydroxide supernatant for 12 hours at 55 degrees C or as recommended by phosphoramidite manufacturer. The ammonium hydroxide solution was evaporated to dryness in a speed-vac and 5'-DMTr-oligonucleotides were purified by HPLC on a C18 reverse-phase matrix using a solvent system of 0.1 M



ammonium acetate and 1:5 ratio 0.1 M ammonium acetate in acetonitrile. Then the oligonucleotides were treated with 80% acetic acid to remove the DMTr group, converted to sodium form and desalted by dialysis against double distilled water. Oligonucleotides were filtered through 0.4 $\mu$  filters, lyophilized and redissolved in double distilled water. Characterization was achieved by denaturing PAGE and MALDI-TOF mass spectrometry.

**Example 2: Synthesis of CpG-PS-oligos containing cytosine analogs**

Following the procedures outlined in Example 1, the following oligonucleotides were synthesized:

Oligo #	Sequence (5'---> 3') and Modification <sup>a</sup>
1	d(CTATCTGACGTTCTCTGT)
2	d(CTATCTGAC <b>*</b> GTTCTCTGT)
3	d(CTATCTGACC <b>*</b> TTCTCTGT)
4	d(CTATCTGAC <b>*</b> GTTCTCTGT)
5	d(CTATCTGACC <b>*</b> TTCTCTGT)

<sup>a</sup> CpG-motif is shown in bold. C\* represents 5-hydroxycytosine (oligos 2 and 3) or N4-ethylcytosine (oligos 4 and 5).

The oligonucleotides were characterized by CGE and MALDI-TOF mass spectrometry (Brucker Proflex III MALDI-TOF mass spectrometer with 337 nm N2 laser). Molecular weights observed and calculated (shown in parentheses) for each oligonucleotide are as follows: Oligo 1, 5704 (5704.8); Oligo 2, 5720 (5720.8); Oligo 3, 5681 (5680.7); Oligo 4, 5733 (5733); Oligo 5, 5694 (5693).

**Example 3: Analysis of spleen weights in treated mice**

Female BALB/c mice (4-5 weeks, 19-21 g, Charles River, Wilmington, MA) were used in the study. The animals were fed with commercial diet and water *ad lib*. The animals were injected intraperitoneally with 5 or 10 mg/kg dose of immunostimulatory oligonucleotide compound dissolved in sterile PBS. One group of mice received PBS alone to serve as a control (PBS). Four animals were used for each immunostimulatory oligonucleotide compound. Mice were sacrificed 72 h later, spleens were harvested and weighed.

**Example 4: Analysis of immunostimulatory oligonucleotide compounds in mouse lymphocyte proliferation assay**

Spleens from CD-1, BALB/c, C57BL/6 mouse (4-8 weeks) were used as source of lymphocytes. Single cell suspensions were prepared by gently mincing with the frosted ends of glass slides. Cells were then cultured in RPMI complete medium [RPMI medium supplemented with 10% fetal bovine serum (FBS) (heat-inactivated at 56 °C for 30 min), 50 µM 2-mercaptoethanol, 100 U/mL penicillin, 100 µg/mL streptomycin, 2 mM L-glutamine]. The cells were then plated in 96-well dishes at a density of 10<sup>6</sup> cells/mL in a final volume of 100 µL. Immunostimulatory oligonucleotide compounds or LPS (lipopolysaccharide) were added to the cell culture in 10 µL of TE buffer (10 mM Tris-HCl, pH 7.5, 1 mM EDTA). The cells were then set to culture at 37 °C. After 44 h, 1 µCi <sup>3</sup>H-uridine (Amersham, Arlington Heights, IL) was added to the culture in 20 µL of RPMI medium, and the cells were pulse-labeled for another 4 h. The cells were harvested by automatic cell harvester (Skatron, Sterling, VA), and the filters were counted by a scintillation counter. The experiments were performed in triplicate.

**Example 5: Lymphocyte proliferatory activity of CpG-PS-oligos containing cytosine analogs**

The immunostimulatory activity of CpG-PS-oligos 1-5 (Example 4) was studied using a BALB/c mouse lymphocyte proliferation assay. In brief, mouse spleen cells were cultured and incubated with CpG-PS-oligos at 0.1, 0.3, 1.0 and 3.0 µg/mL

concentration for 48 hr and cell proliferation was measured by  $^3\text{H}$ -uridine incorporation.

Figure 23 shows the dose-dependent cell proliferatory activity of oligos 1-5 in mouse lymphocyte cultures. At a dose of  $3.0\text{ }\mu\text{g/mL}$ , oligo 1, with natural cytidine, showed a proliferation index of  $29.5\pm 2.1$ . Oligo 2, in which the cytosine base of the deoxycytidine of the CpG-motif is replaced with a 5-hydroxycytosine, also showed dose-dependent lymphocyte proliferation. A proliferation index of  $23.7\pm 2.9$  at  $3.0\text{ }\mu\text{g/mL}$  dose was observed for oligo 2. PS-Oligo 4, which contained N4-ethyl-cytosine in place of the cytosine base in the CpG-motif, also showed dose-dependent cell-proliferation activity. The proliferation index of  $18.7\pm 1.6$  observed for oligo 4 at a dose of  $3\text{ }\mu\text{g/mL}$  suggests that the presence of a bulky hydrophobic substitution on the 4-amino group of cytosine in a CpG-motif slightly impedes immunostimulatory activity.

Oligo 3, in which 5-hydroxy-deoxycytidine was placed in the deoxyguanosine position instead of the deoxycytidine position of the CpG-motif, showed a proliferation index that was similar to that observed for media control (Figure 23). Similarly, the control Oligo 5 in which deoxyguanosine in the CpG-motif was substituted with N4-ethyldeoxycytidine, showed cell proliferation similar to that of media control.

Other oligos, in which cytosine base in the CpG-motif was replaced with 5-methyl-deoxycytosine (2; see Figure 28), 5-methyl-deoxyisocytosine (3), deoxyuridine (5), or deoxy-P-base (7) showed no or insignificant cell proliferatory activity in the same assay system. These results suggest that (i) cell proliferatory activity is maintained when the cytosine base of the CpG motif is replaced with 5-hydroxycytosine or N4-ethylcytosine (Oligos 2 and 4, respectively), but (ii) substitution of the guanine base with these cytosine analogs results in a loss of cell proliferatory activity.

**Example 6: Splenomegaly in mice induced by CpG-PS-oligos containing cytosine analogs**

To confirm the *in vitro* effects of CpG-PS-oligos, Oligos 1, 2, and 4 (from Example 4) were injected intraperitoneally (ip) to BALB/c mice at a dose of 10 mg/kg and the change in spleen weight was measured as an indicator of the level of immunostimulatory activity of each PS-oligo. The change in spleen weight as a result of treatment with CpG-PS-oligos is presented in Figure 24. Female BALB/c mice (4-6 weeks, 19-21 gm) were divided in to different groups with four mice in each group. Oligonucleotides were dissolved in sterile PBS and administered intraperitoneally to mice at a dose of 10 mg/kg. After 72 hr, mice were sacrificed and spleens were harvested and weighed. Each circle represents the spleen weight of an individual mouse and the + represents the mean spleen weight for each group.

Oligo 1, which has natural deoxycytidine in the CpG-motif, showed about 45% increase in spleen weight at a dose of 10 mg/kg, compared with the control group of mice that received PBS. Oligo 2, which has a 5-hydroxycytosine in place of the cytosine base in the CpG-motif, showed about 35% increase in spleen weight at the same dose. Oligo 4, which has N4-ethylcytosine in place of the cytosine base in the CpG-motif, showed about 34% increase in spleen weight at the same dose compared to the control group. These data confirm the results observed in lymphocyte proliferation assays for these oligos containing modified cytidine analogs in place of deoxycytidine in the CpG-motif.

**Example 7: Structure-activity relationships of C\*pG-PS-oligos**

The presence of a methyl group at the 5-position of cytosine (5-methyl-deoxycytosine, 2 (Figure 28)) in a CpG-motif completely abolishes CpG related immunostimulatory effects of CpG-PS-oligos. Based on the results observed in *in vitro* and *in vivo* experiments we have constructed structure-activity relationships for the PS-oligos containing cytosine analogs.

The replacement of the cytosine base (1) in the CpG-motif with 5-methyl-  
isocytosine (3) resulted in complete loss of immunostimulatory activity, as is the case  
with 5-methylcytosine (2), which could be as a result of switching the keto and amino  
groups at the 2 and 4-positions, respectively, and/or placing a hydrophobic methyl  
5 group at the 5-position of cytosine.

Oligo 2, containing a hydrophilic hydroxy substitution at the 5-position of the  
cytosine in the CpG-motif, showed immunostimulatory activity similar to that of oligo  
1, which contains the natural cytosine base. This observation suggests that bulky  
hydrophilic groups are better tolerated than are hydrophobic groups at the 5-position of  
10 cytosine for immunostimulatory activity of CpG-PS-oligos. Perhaps the binding pocket  
for the CpG-oligos on receptor is hydrophilic in nature and can not accommodate a  
hydrophobic group at the 5-position of cytosine.

When the cytosine base in the CpG-motif is replaced with uracil (5 (see Figure  
28)), in which keto groups are present at both the 2 and 4-positions, no  
15 immunostimulatory activity was observed, suggesting that a hydrogen bond donor  
amino group at the 4-position of cytosine is critical for immunostimulatory activity.  
When a large hydrophobic ethyl group is placed on 4-amino group of cytosine in a  
CpG-motif, reduced lymphocyte proliferation and a slightly reduced increase in spleen  
weight in mice were observed, suggesting that a bulky ethyl group at this position does  
20 not interfere with binding of the CpG-PS-oligo to the receptor factors responsible for  
immunostimulatory activity. In spite of the ethyl substitution, the 4-amino group of  
N4-ethylcytosine (6) can participate in hydrogen bond formation with an acceptor. The  
modified pyrimidine base dP, in which the nitrogen group located at the 4-position  
involved in ring structure formation with the 5-position, and which does not have a  
25 hydrogen bond donor amino group at the 4-position, had no mouse lymphocyte  
proliferation activity in cultures, suggesting that the 4-amino group of cytosine in a  
CpG-motif is critical for immunostimulatory activity.

In conclusion, the results presented here show that the functional groups at 2, 3,  
and 4 positions of the cytosine are important for CpG-related immunostimulatory

activity. A hydrophobic substitution at the 5-position of cytosine completely suppresses immunostimulatory activity of a CpG-oligo, while a hydrophilic group at this position is tolerated well. In addition, the immunostimulatory activity of CpG-PS-oligos containing 5-hydroxycytosine or N4-ethylcytosine in place of cytosine in the  
5 CpG-motif can be modulated significantly by incorporating appropriate chemical modifications in the 5'-flanking sequence, suggesting that these cytosine analogs in a CpG-motif are recognized as part of an immunostimulatory motif.

#### Example 8: Synthesis of end-blocked CpG-PS oligonucleotides

The CpG-PS-oligos shown in Figure 17 were synthesized using an automated synthesizer and phosphoramidite approach. Oligo 1 (16-mer) was synthesized using nucleoside-5'- $\beta$ -cyanoethylphosphoramidites. Oligo 2, a 32-mer, was synthesized using nucleoside-3'- $\beta$ -cyanoethylphosphor-amidites and controlled pore glass support (CPG-solid support) with a 3'-linked nucleoside in which 16-mer sequence of Oligo 1 was repeated twice; therefore, Oligo 2 had two 16-mers (Oligo 1) linked by a normal 3'-5'-linkage. Oligo 3, a 32-mer, was synthesized with two 16-mers (Oligo 1) linked by a 5'-5'-linkage, so Oligo 3 had two 3'-ends and no 5'-end. Synthesis of Oligo 3 was carried out in two steps: the first 16-mer was synthesized using nucleoside-3'- $\beta$ -cyanoethylphosphoramidites and solid support with a 3'-linked nucleoside, and then synthesis of the second 16-mer segment was continued using nucleoside-5'- $\beta$ -cyanoethylphosphoramidites. Oligo 4, a 32-mer, comprised two 16-mers (Oligo 1) linked by a 3'-3'-linkage, so Oligo 4 had two 5'-ends and no 3'-end. Synthesis of Oligo 4 was carried out in two steps: the first 16-mer was synthesized using nucleoside-5'- $\beta$ -cyanoethylphosphoramidites and solid support with a 5'-linked nucleoside, and the synthesis of the second 16-mer segment was continued using nucleoside-3'- $\beta$ -cyanoethylphosphoramidites. Synthesis of Oligos 5-8 was carried out by using the same nucleoside- $\beta$ -cyanoethylphosphoramidites as for Oligos 1-4, respectively. At the end of the synthesis, Oligos 1-8 were deprotected with concentrated ammonia solution, purified by reversed phase HPLC, detritylated, desalted and dialyzed. The purity of

each PS-oligo was checked by CGE and the molecular weight was confirmed by MALDI-TOF mass spectral analysis (Table 1). The sequence integrity and directionality of 5'-CpG motif in Oligos 1-8 were confirmed by recording melting temperatures ( $T_m$ s) of the duplexes with their respective DNA complementary strands

- 5 (5'-AAGGTCGAGCGTTCTC-3' for Oligos 1-4, and 5'-ATGGCGCACGCTGGGAGA-3' for Oligos 5-8). The  $T_m$ s of these duplexes were  $53.9 \pm 0.9$  °C (Oligos 1-4),  $61.8$  °C (Oligo 5), and  $58.8 \pm 0.6$  °C (Oligos 6-8) (note that Oligo 5 was a 18-mer and Oligos 6-8 were 32-mers but not 36-mers).

**Example 9: Mouse spleen lymphocyte proliferatory activity of end-blocked CpG-PS oligonucleotides**

Immunostimulatory activity of the end-blocked CpG-PS-oligos of Example 8 was studied initially in a lymphocyte proliferation assay. Typically, mouse (Balb-C) spleen lymphocytes were cultured with CpG-PS-oligos at concentrations of 0.1, 1.0, and 10.0  $\mu$ g/ml for 48 h and cell proliferation was determined by  $^3$ H-uridine incorporation, as described in Example 3. Results are shown in Figure 17

- 20 Oligo 1 induced a dose-dependent effect on cell proliferation; at a concentration of 10  $\mu$ g/ml ( $\sim 2.0$   $\mu$ M), the proliferation index was  $5.0 \pm 0.32$ . Oligo 2, which consisted of two units of Oligo 1 linked by a 3'-5'-linkage, had a proliferation index of  $5.8 \pm 0.28$  at the same dose ( $\sim 1.0$   $\mu$ M). Oligo 3, which consisted of two units of Oligo 1 linked by a 5'-5'-linkage, had a proliferation index of  $2.0 \pm 0.26$ , reflecting a significantly lower immunostimulatory activity than observed with Oligos 1 and 2. Oligo 4, which consisted of two units of Oligo 1 linked by a 3'-3'-linkage, had a proliferation index of  $7.2 \pm 0.5$ , reflecting a greater immunostimulatory activity than observed with Oligos 1 and 2.

- 25 Similar results were obtained with Oligos 5-8. Oligo 5 had a proliferation index of  $3.9 \pm 0.12$ . Oligos 6-8, in which two units of Oligo 5 are linked by a 3'-5'-linkage (Oligo 6), 5'-5'-linkage (Oligo 7), and 3'-3'-linkage (Oligo 8) had proliferation indices of  $4.9 \pm 0.2$ ,  $1.74 \pm 0.21$ , and  $7.7 \pm 0.82$ , respectively. Comparison of the results obtained

with Oligos 6-8 show that Oligos 6 and 8, in which two Oligo 5 sequences were linked by a 3'-5'-linkage or a 3'-3'-linkage had greater immunostimulatory activity, while Oligo 7, in which two Oligo 5 were linked by a 5'-5'-linkage had significant less immunostimulatory activity, than did Oligo 5.

5       Based on lymphocyte proliferation results of Oligos 1-8, it is clear that when oligos are linked through their 5'-ends, there is a significant loss of immunostimulatory activity, while if they are linked through their 3'-ends, there is an increase in immunostimulatory activity. It is important to note that 3'-3'-linked oligos have shown substantially greater stability towards degradation by exonucleases than the oligos that contained a free 3'-end, which could also result in increased immunostimulatory activity. The lower immunostimulatory activity of Oligos 3 and 7, in which the 5'-end of oligos is blocked, suggests that accessibility to 5'-end of oligo is essential for immunostimulatory activity of CpG-PS-oligos.

10  
13  
16  
19  
22  
25  
28  
31  
34  
37  
40  
43  
46  
49  
52  
55  
58  
61  
64  
67  
70  
73  
76  
79  
82  
85  
88  
91  
94  
97  
100  
103  
106  
109  
112  
115  
118  
121  
124  
127  
130  
133  
136  
139  
142  
145  
148  
151  
154  
157  
160  
163  
166  
169  
172  
175  
178  
181  
184  
187  
190  
193  
196  
199  
202  
205  
208  
211  
214  
217  
220  
223  
226  
229  
232  
235  
238  
241  
244  
247  
250  
253  
256  
259  
262  
265  
268  
271  
274  
277  
280  
283  
286  
289  
292  
295  
298  
301  
304  
307  
310  
313  
316  
319  
322  
325  
328  
331  
334  
337  
340  
343  
346  
349  
352  
355  
358  
361  
364  
367  
370  
373  
376  
379  
382  
385  
388  
391  
394  
397  
400  
403  
406  
409  
412  
415  
418  
421  
424  
427  
430  
433  
436  
439  
442  
445  
448  
451  
454  
457  
460  
463  
466  
469  
472  
475  
478  
481  
484  
487  
490  
493  
496  
499  
502  
505  
508  
511  
514  
517  
520  
523  
526  
529  
532  
535  
538  
541  
544  
547  
550  
553  
556  
559  
562  
565  
568  
571  
574  
577  
580  
583  
586  
589  
592  
595  
598  
601  
604  
607  
610  
613  
616  
619  
622  
625  
628  
631  
634  
637  
640  
643  
646  
649  
652  
655  
658  
661  
664  
667  
670  
673  
676  
679  
682  
685  
688  
691  
694  
697  
700  
703  
706  
709  
712  
715  
718  
721  
724  
727  
730  
733  
736  
739  
742  
745  
748  
751  
754  
757  
760  
763  
766  
769  
772  
775  
778  
781  
784  
787  
790  
793  
796  
799  
802  
805  
808  
811  
814  
817  
820  
823  
826  
829  
832  
835  
838  
841  
844  
847  
850  
853  
856  
859  
862  
865  
868  
871  
874  
877  
880  
883  
886  
889  
892  
895  
898  
901  
904  
907  
910  
913  
916  
919  
922  
925  
928  
931  
934  
937  
940  
943  
946  
949  
952  
955  
958  
961  
964  
967  
970  
973  
976  
979  
982  
985  
988  
991  
994  
997  
1000

**Example 10: Splenomegaly in mice induced by end-blocked CpG-PS oligonucleotides**

To confirm the immunostimulatory activity of Oligos 1-8 (Example 8) *in vivo*, a dose of 5 mg/kg of oligonucleotides was injected intraperitoneally to Balb-C mice. The mice were sacrificed 72 hours post-administration, spleens were removed, blotted to dryness, and weighed. Change in spleen weight in treated and untreated mice was used as a parameter for immunostimulatory activity.

Administration 5 mg/kg dose of Oligo 1 caused about 40% increase in spleen weight compared with the control mice that received PBS. Administration of Oligos 2 and 4 also caused about 50% increase in spleen weight. Administration of Oligo 3 caused no difference in spleen weight compared with control mice. These results further support the observation that Oligo 3, in which 5'-end was blocked, had significantly less immunostimulatory activity compared to oligos that had accessible 5'-end. These results were also confirmed with the administration of Oligos 5-8. Administration of Oligos 5, 6, and 8 caused about 40-50% increase in spleen weight,



whereas no change in spleen weight was observed following the administration of Oligo 7.

The above results suggest that the immunostimulatory activity of PS-oligos containing a CpG motif is significantly minimized if the 5'-end of the oligo is not accessible. This loss in immunostimulatory activity of Oligos 3 and 7 cannot be explained based on nuclease stability, as both oligos have two 3'-ends and are not more susceptible to 3'-exonuclease degradation than are Oligos 1, 2, 5, and 6, which have one 3'-end. PS-Oligos 4 and 8, which have their 3'-ends blocked and are very stable to degradation by exonucleases, showed similar immunostimulatory activity. Oligos 4 and 8 may show sustained immunostimulatory activity due to their increased *in vivo* stability, which is not evident in the present study as mice were sacrificed at only 72 hours after administration. Studies are in progress in which mice will be sacrificed at times later than 72 hours after administration.

The results described here are intriguing and suggest that the 5'-end of CpG-PS-oligos is critical for immunostimulatory activity. As discussed here, we have shown that substitution of deoxynucleosides in 5'-flanking regions by modified 2'- or 3'-substituted ribonucleosides resulted in increased immunostimulatory activity. In addition, substitution of deoxynucleosides immediately upstream (5'-end) to the CpG motif caused a significant suppression and substitution of deoxynucleosides immediately downstream (3'-end) to the CpG motif had no effect on immunostimulatory activity. Taken together, these results suggest that the enzyme/receptor responsible for the immunestimulation recognizes the CpG motif in oligos from the 5'-end and requires accessibility to the 5'-end.

While the foregoing invention has been described in some detail for purposes of clarity and understanding, it will be appreciated by one skilled in the art from a reading of this disclosure that various changes in form and detail can be made without departing from the true scope of the invention and appended claims.